

β -NAD⁺-Li


β -nicotinamide-adenine dinucleotide (= β -NAD), oxidized form (mono lithium salt)

β -Diphosphopyridine (= β -DPN), oxidized form (mono lithium salt)

Coenzyme-I, oxidized form (mono lithium salt)

from Yeast

Structure



Formula : $C_{21}H_{27}N_7O_{14}P_2 \cdot Li$

Formula weight : 669.4

Specification

Purity

Determined by Enzymatic Method (ADH)

Water Content

Li

UV Spectral Analysis

ϵ at 260 nm and pH 7.5

Ratio at pH 10

$$A_{250}/A_{260}$$

$$A_{280}/A_{260}$$

ϵ when reduced with ADH at 340 nm and pH 10

Ratio when reduced with ADH at pH 10

$$A_{340}/A_{260}$$

Specifications

$\geq 95\%$

$< 8\%$

$1.0 \pm 0.5\%$

$(18.5 \pm 0.5) \times 10^3$

0.83 ± 0.03

0.21 ± 0.02

$(6.3 \pm 0.2) \times 10^3$

0.43 ± 0.01

Assay Procedure

I. Spectrophotometric Method

Wavelength ; 340 nm, Light path length ; 1 cm

Pipette the following reagents into a cuvette

	a	b	c
Tris-EtOH (0.1 mol/L, 2.4%)	5.0 mL	5.0 mL	5.0 mL
ADH (50 IU/mL)	0.3 mL	—	0.3 mL
NAD ⁺ (0.45 mg/mL)	0.5 mL	0.5 mL	—
Distilled water	0.2 mL	0.5 mL	0.7 mL

II. Calculation

$$\frac{\Delta A \cdot V \cdot MW \times 100}{6.3 \times 10^3 \cdot d \cdot v \cdot s} \times \frac{100}{100 - W - Li} = \text{Purity of NAD}^+$$

$\Delta A = A_a - (A_b + A_c)$

V = Total volume of reaction mixture (6.0 mL)

MW = 663.4, as of anhydrate

6.3×10^3 = Molar extinction coefficient of NADH at 340 nm ($L \cdot mol^{-1} \cdot cm^{-1}$)

d = Light path length (1 cm)


v = Sample volume (0.5 mL)

s = Sample concentration (0.45 mg/mL)

W = Water Content (%)

Li = Li (%)

Reference Data



Storage

Keep tightly stoppered in the dark below 5°C.

For prolonged storage keep below -20°C.

OYC No./Package

OYC No. 44097900
Package Bulk

(Research reagent use only, not for medical use.)



ORIENTAL YEAST CO.,LTD.