

# rLDH(CH)

**recombinant Lactate dehydrogenase EC 1.1.1.27**

*from Chicken heart*

## Reaction Equation



## Specification

### Specific Activity

U/mg protein > 200 units

### Contaminants

Malate dehydrogenase	< 0.03%
Myokinase	< 0.01%
Pyruvate kinase	< 0.003%
Glutamic-pyruvic transaminase*	< 0.03%
Glutamic-oxaloacetic transaminase*	< 0.03%

\*Including  $\alpha$ -Hydroxyglutarate dehydrogenase activity

## Properties

pH stability : pH 6.3 - 9.2 (25°C, 1 week)

Thermal stability :  $\leq 65^\circ\text{C}$  (pH 7.5, 10 min)

Optimum pH : 7.0 - 7.5

Optimum temp. :  $\geq 37^\circ\text{C}$

Km value :  $4.1 \times 10^{-5}$  mol/L (Pyruvate)

$1.8 \times 10^{-5}$  mol/L (NADH)

Molecular weight : 36 kDa (SDS-PAGE)

## Assay Procedure

### I Spectrophotometric Method

Wavelength : 340 nm, Light path length : 1 cm

Final volume : 3.17 mL, Temperature : 25°C

Pipette the following reagents into a cuvette

3.00 mL	K-phosphate buffer (0.1 mol/L, pH 7.0)
0.10 mL	Li-pyruvate or Na-pyruvate (25.4 mmol/L)
0.05 mL	NADH (10 mg/mL) dissolved in Tris (10 mmol/L)
0.02 mL	rLDH (CH) (approx. 3 U/mL)

## II Calculation

$$\frac{\Delta A/\text{min} \cdot V \cdot D}{6.3 \cdot d \cdot v} = \text{U/mL}$$

$\Delta A/\text{min}$  = The change in absorbance at 340 nm/minute

V = Total volume of reaction mixture (3.17 mL)


D = Enzyme dilution factor

6.3 = mmol/L extinction coefficient of NADH  
( $\text{L} \cdot \text{mmol}^{-1} \cdot \text{cm}^{-1}$ )

d = Light path length (1 cm)

v = Volume of enzyme sample (0.02 mL)

## Reference Data



## Preparation and Storage

Lyophilized powder

Store below -20°C

## Cat. No./Package

Cat. No.	Package
46757903	Bulk

For in vitro diagnostic or research use only



ORIENTAL YEAST CO.,LTD.